

Projective Properties Of Special (A, B)- Finsler Metrics On Γ – Curvature Tensor Of C3 – Like Conformal Finsler Space

Gayathri. K¹ and Narasimhamurthy. S. K²

¹Assistant Professor, Department of Mathematics, S.I.T, Tumkur-572103, India

²Professor and Chairman, Department of Mathematics,
Kuvempu University, Shankaragatta, India.

Abstract: The purpose of the present paper is to study the properties of the γ – curvature tensor in C3 – like Finsler space F^n of dimension ($n \geq 4$), in which the conformal Cartan torsion tensor \overline{C}_{ijk} is said to be a conformal C3 – like Finsler space. Finsler geometry is originated from Differential geometry. Finsler geometry is Riemannian metric without quadratic restriction. In Finsler space we see special metrics such as Randers metric, Kropina metric and Matsumoto metric.,etc. Projective change between two Finsler metrics arise from Information Geometry. Such metrics have special geometric properties and will play an important role in Finsler geometry. In this paper, we are going to study class of Projective change between two (α, β) – metrics, which are defined as the sum of a Riemannian metric and 1 – form.

Keywords: Finsler metric, C-reducible, C3-like, γ - curvature tensor, Special Finsler metric, (α, β) – metric, Douglas Space, Geodesic, Spray coefficients, Projectively related metric, Projective change between two metrics.

I. PRELIMINARIES

Let $F^n = (M^n, L)$ be a Finsler space on a differential manifold M endowed with a fundamental function $L(x, y)$.

We use the following notations [2], [6]:

- $g_{ij} = \frac{1}{2} \dot{\partial}_i \dot{\partial}_j L^2, \quad g^{ij} = (g_{ij})^{-1}, \quad \dot{\partial}_i = \frac{\partial}{\partial y^i},$
- $C_{ijk} = \frac{1}{2} \dot{\partial}_k g_{ij}, \quad C_{ij}^k = \frac{1}{2} g^{km} (\dot{\partial}_m g_{ij}),$
- $h_{ij} = g_{ij} - l_i l_j, \quad h_k^i = \delta_k^i - l^i l_k,$
- $m_i = b_i - \beta L^{-1} l^i,$
- $C_{ij}^h l_h = 0,$
- $h_k^i m_i = m_k,$

$$g) \quad l^i m_i = 0,$$

Where l_i, m_i and n_i are the unit vectors and h_{ij} is a angular metric tensor.

An interesting result concerned with the theory of projective change was given by Rapsca's paper. He proved necessary and sufficient conditions for projective change.

S. Bacso and M. Matsumoto [2] discussed the projective change between Finsler spaces with (α, β) – metric.

H. S. Park and Y. Lee [6] studied on projective changes between a Finsler space with (α, β) – metric and the associated Riemannian metric.

X.Cheng and Z. Shen studied a class of Finsler metrics with isotropic S-curvature and curvature properties of Finsler metrics.

Recently some results on a class of (α, β) – metrics with constant flag curvature have been studied by Ningwei Cui, Yi-Bing Shen [5], N. Cui and Z. Lin.

II. SOME IMPORTANT DEFINITIONS

2.1 Definition:

Let $F^n = (M^n, L(x, y))$ and $\bar{F}^n = (M^n, \bar{L}(x, y))$ be two Finsler spaces on the same underlying manifold M^n . If the angle in F^n is equal to that in \bar{F}^n for any tangent vectors, then F^n is called conformal to \bar{F}^n and the change $L \rightarrow \bar{L} = e^\sigma L$ of the metric is called a conformal change and $\sigma(x)$ is a conformal factor.

Example 1: We consider a Finsler space $F^n = (M^n, L(\alpha, \beta))$, where α is a Riemannian metric, β is a 1-form and a conformal change $L(\alpha, \beta) \rightarrow \bar{L} =$

$e^{\sigma(x)}L(\alpha, \beta)$. Since $L(\alpha, \beta)$ is assumed to be (1)*p* – homogeneous in α and β , we get $\bar{L} = L(\bar{\alpha}, \bar{\beta})$, where $\bar{\alpha} = e^{\sigma(x)}\alpha$ and $\bar{\beta} = e^{\sigma(x)}\beta$. Thus the conformal change gives rise to the change $(\alpha, \beta) \rightarrow (\bar{\alpha}, \bar{\beta}) = (e^{\sigma(x)}\alpha, e^{\sigma(x)}\beta)$ of the pair (α, β) independently of the form of the function $L(\alpha, \beta)$. Thus we also get the conformal change $\alpha \rightarrow \bar{\alpha} = e^{\sigma(x)}\alpha$ of the associated Riemannian space $R^n = (M^n, \alpha)$.

Under the conformal change, we get the following relations [3], [4]:

- a) $\bar{g}_{ij} = e^{2\sigma}g_{ij}$, $\bar{g}^{ij} = e^{-2\sigma}g^{ij}$,
- b) $\bar{C}_{ijk} = e^{2\sigma}C_{ijk}$,
 $\bar{C}_{jk}^i = C_{jk}^i$, $\bar{C}_k^i = \bar{C}_k = C_{jk}^i = C_k$,
- c) $\bar{l}^i = e^{-\sigma}l^i$, $\bar{l}_i = e^{\sigma}l_i$, $\bar{y}_i = e^{2\sigma}y_i$, (2.1)
- d) $\bar{h}_{ij} = e^{2\sigma}h_{ij}$, $\bar{h}_j^i = h_j^i$,
- e) $\bar{L} = e^{\sigma}L$,
- f) $\bar{m}_k = e^{\sigma}m_k$.

2.2 Definition: A Finsler space is said to be C-reducible if it satisfies the equation $C_{ijk} = \frac{C_i h_{jk} + C_j h_{ki} + C_k h_{ij}}{n+1}$

2.3 Definition: A Finsler metric is a scalar field $L(x, y)$ which satisfies the following three conditions:

- i. It is defined and differentiable at any point of $TM^n \setminus \{0\}$,
- ii. It is positively homogeneous of first degree in y^i , that is, $L(x, \lambda y) = \lambda L(x, y)$, for any positive number λ ,
- iii. It is regular, that is, $g_{ij}(x, y) = \frac{1}{2} \dot{\partial}_i \dot{\partial}_j L^2$, constitute the regular matrix g_{ij} , where $\dot{\partial}_i = \frac{\partial}{\partial y^i}$.

The manifold M^n equipped with a fundamental function $L(x, y)$ is called Finsler metric $F^n = (M^n, L)$.

2.4 Definition: Two Finsler metrics L and \bar{L} are projectively related if and only if their spray coefficients have the relation

$$G^i = \bar{G}^i + P(y)y^i \quad (2.3)$$

2.5 Definition: A Finsler metric is projectively related to another metric if they have the same

geodesics as point sets. In Riemannian geometry, two Riemannian metrics α and $\bar{\alpha}$ are projectively related if and only if their spray coefficients have the relation

$$G_{\alpha}^i = G_{\bar{\alpha}}^i + \lambda_{x^k} y^k y^i \quad (2.4)$$

2.6 Definition: Let

$$\phi(s) - s\phi'(s) + (b^2 - s^2)\phi''(s) > 0, \quad (|s| \leq b \leq b_0). \quad (2.5)$$

If $\alpha = \sqrt{a_{ij}y^i y^j}$ is a Riemannian metric and $\beta = b_i y^i$ is 1-form satisfying $\|\beta_x\|_{\alpha} < b_0 \quad \forall x \in M$, then $L = \phi(s)$, $s = \beta/\alpha$, is called an (regular) (α, β) – metric. In this case, the fundamental form of the metric tensor induced by L is positive definite.

2.7 Definition: Let $F^n = (M^n, L)$ and $\bar{F}^n = (M^n, \bar{L})$ be two Finsler spaces on a common underlying manifold M^n . If any geodesic on F^n is also a geodesic on \bar{F}^n and the converse is true, then the change $L \rightarrow \bar{L}$ of the metric is called a projective change.

(2.2)

The relation between the geodesic coefficients G^i of L and geodesic coefficients G_{α}^i of α is given by

$$G^i = G_{\alpha}^i + \alpha Q s_0^i + \{-2Q\alpha s_0 + r_{00}\} \{\Psi b^i + \Theta \alpha^{-1} y^i\}, \quad (2.6) \quad \text{where,}$$

$$\Theta = \frac{\phi \phi' - s(\phi \phi'' + \phi' \phi')}{2\phi((\phi - s\phi') + (b^2 - s^2)\phi'')}, \quad Q = \frac{\phi'}{\phi - s\phi'},$$

$$\Psi = \frac{1}{2} \frac{\phi''}{(\phi - s\phi') + (b^2 - s^2)\phi''}$$

2.8 Definition: Let $D_{jkl}^i = \frac{\partial^3}{\partial y^j \partial y^k \partial y^l} (G^i - \frac{1}{n+1} \frac{\partial G^m}{\partial y^m} y^i)$, where G^i are the spray coefficients of L . The tensor $D = D_{jkl}^i \partial_i \otimes dx^j \otimes dx^k \otimes dx^l$ is called the Douglas tensor. A Finsler metric is called Douglas metric if the Douglas tensor vanishes.

Then there exists a class of scalar functions $H_{jk}^i = H_{jk}^i(x)$, such that

$$H_{00}^i = T^i - \bar{T}^i \frac{1}{n+1} (T_{y^m}^m - \bar{T}_{y^m}^m) y^i, \quad (2.7)$$

Where $H_{00}^i = H_{jk}^i y^j y^k$, T^i and $T_{y^m}^m$ are given by the relations

$$T^i = \alpha Q s_0^i + \Psi \{-2Q\alpha s_0 + r_{00}\} b^i \quad (2.8)$$

and

$$T_{y^m}^m = Q' s_0 + \Psi' \alpha^{-1} (b^2 - s^2) [r_{00} - 2Q\alpha s_0] + 2\Psi [r_0 - Q'(b^2 - s^2) s_0 - Q s s_0] \quad (2.9)$$

I. Properties of C3-like conformal Finsler space:

3.1 Theorem : A C3-like conformal Finsler space reduce to a semi-C-reducible conformal Finsler space if the vectors a_i and b_i are parallel to C_i .

II. The γ -curvature tensor of C-reducible Finsler space:

3.2 Theorem : The C-reducible Finsler space and γ -curvature tensor S_{hijk} satisfies the symmetric property and it satisfies $C_k y_j = C_j y_k$.

Corollary 3.1: Under conformal change, the C-reducible condition and ϑ -curvature tensor also satisfy property $C_k y_j = C_j y_k$.

Example 2: Let T_{ij} be a tensor of (0,2)-type of a two dimensional Finsler space and $T_{\alpha\beta}$ be scalar components of T_{ij} with respect to the Berwald frame:

$$T_{ij} = T_{11} l_i l_j + T_{12} l_i m_j + T_{21} m_i l_j + T_{22} m_i m_j$$

If T_{ij} is symmetric, we have $T_{12} = T_{21}$, and if T_{ij} is skew-symmetric, then $T_{0j} = 0$, $T_{ij} = 0$; therefore, by this condition, the ϑ -curvature tensor S_{hijk} of CT of any two dimensional Finsler space vanishes identically.

Projective change between two Finsler metrics

: In this section, we find the projective relation between two (α, β) - metrics, that is, Special (α, β) - metric $L = 1 + c_c \beta + \frac{\beta^2}{\alpha}$ and Randers metric $\bar{L} = \bar{\alpha} + \bar{\beta}$ on a same underlying manifold M of dimension $n > 2$.

From (2.3), $L = 1 + c_c \beta + \frac{\beta^2}{\alpha}$ is a regular Finsler metric if and only if 1-form β satisfies the condition $\|\beta_x\|_\alpha < \frac{1}{2}$ for any $x \in M$. The geodesic coefficients are given by (2.4) with ,

$$\theta = \frac{c_c - 4c_c s + c_c s^2 - 4s^3}{1 + 2b^2 + 2c_c b^2 s + 2s^2(b^2 - 1) - 3c_c s^3 - 3s^4}, \quad Q = \frac{c_c + 2s}{1 - s^2}, \quad \Psi = \frac{1}{(1 + 2b^2) - 3s^2} \quad (3.1)$$

Substituting (3.1) into (2.4), we get

$$G^i = G_\alpha^i + \frac{\alpha^2 (c_c \alpha + 2\beta)}{\alpha^2 - \beta^2} s_0^i + \left\{ \frac{-2\alpha^2 (c_c \alpha + 2\beta)}{\alpha^2 - \beta^2} s_0 + r_{00} \right\} \left\{ \frac{\alpha^2}{(1 + 2b^2)\alpha^2 - 3\beta^2} b^i + \frac{y^i}{(1 + 2b^2)\alpha^4 + 2c_c b^2 \alpha^3 \beta + 2(b^2 - 1)\alpha^2 \beta^2 - 3c_c \alpha \beta^3 - 3\beta^4} \right\}$$

From (2.3), $\bar{L} = \bar{\alpha} + \bar{\beta}$ is a regular Finsler metric if and only if $\|\beta_x\|_\alpha < 1$ for any $x \in M$.

The geodesic coefficients are given by (2.4) with

$$\bar{\theta} = \frac{1}{2(1+s)}, \quad \bar{Q} = \mathbf{1}, \quad \bar{\Psi} = \mathbf{0} \quad (3.3)$$

First we prove the following lemma:

Lemma 3.1: Let $L = 1 + c_c\beta + \frac{\beta^2}{\alpha}$ and $\bar{L} = \bar{\alpha} + \bar{\beta}$ be two (α, β) – metrics on a manifold M with dimension $n > 2$. Then they have the same Douglas tensor if and if both the metrics L and \bar{L} are Douglas metrics.

Proof: First, we prove the sufficient condition. Let L and \bar{L} be Douglas metrics and corresponding Douglas tensors be D_{jkl}^i and \bar{D}_{jkl}^i . Then by the definition of Douglas metric, we have $D_{jkl}^i = 0$ and $\bar{D}_{jkl}^i = 0$, that is, both L and \bar{L} have same Douglas tensor. Next, we prove the necessary condition. If L and \bar{L} have the same Douglas tensor, then (2.5) holds. Substituting (3.1) and (3.3) in to (2.5), we obtain

$$H_{00}^i = \frac{A^i\alpha^{10} + B^i\alpha^8 + C^i\alpha^6 + D^i\alpha^4 + E^i\alpha^2}{I\alpha^8 + J\alpha^6 + K\alpha^4 - J\alpha^2 + L} - \bar{\alpha}\bar{s}_0^i \quad (3.4)$$

Where $A^i = (n+1)(1+2b^2)\{c_c(1+2b^2)s_0^i - 2c_c s_0 b^i\}$,

$$B^i = (n+1)\{(1+2b^2)[2(1+2b^2)\beta s_0^i - 4s_0 b^i \beta - 3c_c \beta^2 s_0^i] - 2(2+b^2)[c_c(1+2b^2)s_0^i - 2c_c s_0 b^i] - 4(1+2b^2)^2 s_0\},$$

$$C^i = (n+1)\{3[(1+2b^2)s_0^i - 2s_0 b^i] + 6(2+b^2)s_0^i c_c \beta^4 - [6s_0^i + 4(2+b^2)(1+2b^2)s_0^i - 4s_0 b^i]\beta^3 - 4[(2+b^2)(1+2b^2)s_0]\beta^2 + r_{00} b^i\},$$

$$D^i = (n+1)\{12(2+b^2)s_0^i r_{00} b^i \beta^5 + 3[c_c(1+2b^2)s_0^i - 2c_c s_0 b^i] + 6s_0(1+2b^2)\beta^4 - 2(2+b^2)r_{00} b^i \beta^2\},$$

$$E^i = (n+1)\{3\beta^4 r_{00} b^i - 12s_0^i \beta^7\}$$

$$\lambda = \frac{1}{n+1}$$

$$\text{and } I = (1+n)(1+2b^2)^2,$$

$$J = 2[4+4b^4+11b^2]\beta^2,$$

$$K = (n+1)[3+58b^2+4b^4]\beta^4,$$

$$L = -12(n+1)[2+b^2]\beta^6,$$

$$M = 9\beta^4 \quad (3.6)$$

Then (3.4) equivalent to

$$A^i\alpha^{10} + B^i\alpha^8 + C^i\alpha^6 + D^i\alpha^4 + E^i\alpha^2 = (I\alpha^8 + J\alpha^6 + K\alpha^4 + L\alpha^2 + M)(H_{00}^i + \bar{\alpha}\bar{s}_0^i) \quad (3.7)$$

Replacing y^i by $-y^i$ in (3.7) yields

$$-A^i\alpha^{10} + B^i\alpha^8 - C^i\alpha^6 + D^i\alpha^4 - E^i\alpha^2 = (I\alpha^8 - J\alpha^6 + K\alpha^4 - L\alpha^2 + M)(H_{00}^i - \bar{\alpha}\bar{s}_0^i) \quad (3.8)$$

Subtracting (3.8) from (3.7), we obtain

$$A^i\alpha^{10} + C^i\alpha^6 + E^i\alpha^2 = H_{00}^i\alpha^2(I\alpha^6 + J\alpha^4 + K\alpha^2 + L) + \alpha\bar{\alpha}\bar{s}_0^i(M) \quad (3.9)$$

Now, we study two cases for Riemannian metric.

Case (i): If $\bar{\alpha} = \mu(x)\alpha$, then (3.9) reduces to

$$A^i\alpha^{10} + C^i\alpha^6 + E^i\alpha^2 = H_{00}^i\alpha^2(I\alpha^6 + J\alpha^4 + K\alpha^2 + L) + \mu(x)\alpha^2\bar{s}_0^i(M) \quad (3.10)$$

above equation can be written as

$$H^i = [H_{00}^i(I\alpha^6 + J\alpha^4 + K\alpha^2 + L) + \mu(x)\bar{s}_0^i(K) - A^i\alpha^8 - C^i\alpha^4 - E^i]\alpha^2 \quad (3.11)$$

From (3.11), we can observe H^i has the factor α^2 , that is, $12\lambda y^i r_{00} \beta^4$ has the factor α^2 . Since β^2 is not having α^2 factor, the only possibility is that βr_{00} has the factor α^2 . Then for each i there exists a scalar function $\tau^i = \tau(x)$ such that $\beta r_{00} = \tau^i \alpha^2$,

$$\text{which is equivalent to } b_j r_{0k} + b_k r_{0j} = 2\tau^i \alpha_{jk} .$$

If $n > 2$ and assuming $\tau^i \neq 0$, then $2 \geq \text{rank}(b_j r_{0k}) + \text{rank}(b_k r_{0j})$

$$\begin{aligned} &> \\ \text{rank}(b_j r_{0k} + b_k r_{0j}) & \\ &= \\ \text{rank}(2\tau^i \alpha_{jk}) &> 2 , \end{aligned}$$

which is impossible unless $\tau^i = 0$. Then $\beta r_{00} = 0$. Since $\beta \neq 0$, we have $r_{00} = 0$, implies that $b_{ij} = 0$.

Case (ii): If $\bar{\alpha} \neq \mu(x)\alpha$, then from (3.9), observe H^i has the factor α , that is, $12\lambda y^i \beta^4 r_{00}$ has the factor α . Note that β^2 has no factor α . Then the only possibility is that βr_{00} has the factor α^2 . As in the case(i), we have $b_{ij} = 0$ when $n > 2$.

Special (α, β) - metric is a Douglas metric if and only if $b_{ij} = 0$. Thus L is a Douglas metric. Since L is projectively related to \bar{L} , then both L and \bar{L} are Douglas metrics.

Now, we prove the following theorem:

Theorem 3.1: The Finsler metric $L = 1 + c_c \beta + \frac{\beta^2}{\alpha}$ is projectively related to if and only if the following conditions are satisfied

$$G_\alpha^i = G_\alpha^i + P y^i , \quad b_{ij} = 0 , \quad d\bar{\beta} = 0, \quad (3.12)$$

Where $b = \|\beta\|_\alpha$, b_{ij} denote the coefficients of the covariant derivatives of β with respect to α , P is a scalar function.

Proof: Let us prove the necessary condition. Since Douglas tensor is an invariant under projective changes between two Finsler metrics, if L is projectively related to \bar{L} , then they have the same Douglas tensor. According to lemma 3.1, we get both L and \bar{L} are Douglas metrics.

Since Randers metric $\bar{L} = \bar{\alpha} + \bar{\beta}$ is a Douglas metric if and only if $\bar{\beta}$ is closed, we have,

$$d\bar{\beta} = 0 \quad (3.13)$$

and $L = 1 + c_c \beta + \frac{\beta^2}{\alpha}$ is a Douglas metric if and only if

$$b_{ij} = 0 , \quad (3.14)$$

Where b_{ij} denote the coefficients of the covariant derivatives of $\beta = b_i y^i$ with respect to α . In this case, β is closed. Hence $s_{ij} = 0$, implies that $b_{ij} = b_{j|i}$. Thus $s_0^i = 0$, $s_0 = 0$.

By using (3.14), we have $r_{00} = r_{ij} y^i y^j = 0$. Substituting all these in (3.2), we obtain

$$G^i = G_\alpha^i \quad (3.15)$$

Since L is projective to \bar{L} , this is a Randers change between L and $\bar{\alpha}$. Since $\bar{\beta}$ is closed, then L is projectively related to $\bar{\alpha}$. Thus there is a scalar function $P = P(y)$ on $TM \setminus \{0\}$ such that

$$G^i = G_{\bar{\alpha}}^i + Py^i \quad (3.16)$$

From (3.15) and (3.16), we have

$$G_{\alpha}^i = G_{\bar{\alpha}}^i + Py^i \quad (3.17)$$

Equations (3.13) and (3.14) together with (3.17) complete the proof of the necessary condition.

Since $\bar{\beta}$ is closed, it is sufficient to prove that L is projectively related to $\bar{\alpha}$. Substituting (3.14) in to (3.2) yields (3.15).

From (3.15) and (3.17), we have $G^i = G_{\bar{\alpha}}^i + Py^i$

That is, L is projectively related to $\bar{\alpha}$.

From the previous theorem, we get the following corollaries.

Corollary 3.2: The Finsler metric $L = \alpha + \frac{\beta^2}{\alpha}$ is projectively related to $\bar{L} = \bar{\alpha} + \bar{\beta}$ if and only if they are Douglas metrics and the spray coefficients of α and $\bar{\alpha}$ have the following relation

$$G_{\alpha}^i = G_{\bar{\alpha}}^i + Py^i$$

Where P is a scalar function.

In this, we assume that the Randers metric $\bar{L} = \bar{\alpha} + \bar{\beta}$ is locally Minkowskian, where $\bar{\alpha}$ is an Euclidean metric and $\bar{\beta} = \bar{b}_i y^i$ is a one form with $\bar{b}_i = \text{constants}$. Then (3.12) can be written as

$$G_{\alpha}^i = Py^i, \quad b_{ij} = 0 \quad (3.18)$$

Thus, we state

Corollary 3.3: The Finsler metric $L = 1 + c_c \beta + \frac{\beta^2}{\alpha}$ is projectively related to $\bar{L} = \bar{\alpha} + \bar{\beta}$ if and only if L is projectively flat, in other words, L is projectively flat if and only if (3.18) holds.

CONCLUSION

- The conformal γ -curvature tensor \bar{S}_{hijk} on a C3-like conformal Finsler space reduces to the conformal change condition.
- The Finsler metric $L = 1 + c_c \beta + \frac{\beta^2}{\alpha}$ is projectively related to $\bar{L} = \bar{\alpha} + \bar{\beta}$ if and only if L is projectively flat, in other words, L is projectively flat if and only if $G_{\alpha}^i = Py^i$, $b_{ij} = 0$ holds true.

REFERENCES

- [1] P.L. Antonelli, R.S Ingarden and M. Matsumoto, *The Theory of spray and Finsler spaces with applications in Physics and Biology*, Kluwer academic publishers, London, 1985.
- [2] S. Bacso and M. Matsumoto, *Projective change between Finsler spaces with (α, β) -metric*, Tensor N. S., 55(1994), 252-257.
- [3] S. Bacso, X. Cheng and Z. Shen, *Curvature Properties of (α, β) -metrics*, Adv. Stud. Pure Math. Soc., Japan (2007).
- [4] Chern S.S and Z. Shen, *Riemann-Finsler Geometry*, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2005.

- [5] X.Cheng and Z. Shen, *A Class of Finsler metrics with isotropic S-curvature*, Israel J. Math., 169(2009), 317-340.
- [6] Kitayama M., *Finsler spaces admitting a parallel vector field*, Balkan J. Geometry and its Appl., **3** (1998), 29-36.
- [7] Matsumoto M., *Foundations of Finsler geometry and Special Finsler spaces*, Kaiseisha press, Otsu, Saikawa, 1986.
- [8] M. Matsumoto and X. Wei, *Projective changes of Finsler spaces of constant curvature*, Publ. Math. Debreen, 44(1994), 175-181.
- [9] B. Nafaji, Z.Shen and A. Tayebi, *Finsler metrics of scalar flag curvature with special non-Riemannian curvature properties*, J.Gem. Dedicata., 131 (2008), 87-97.
- [10] Narasimhanurthy S.K., Bagewadi C.S. and Nagaraja H.G., *Infinitesimal C-conformal motions of special Finsler spaces*, Tensor, N.S., **64** (2003), 241-247.
- [11] Narasimhanurthy S.K and Bagewadi C.S., *C-conformal special Finsler spaces admitting a parallel vector field*, **65** (2004), 162-169.
- [12] Ningwei Cui and Yi-Bing Shen, *Projective change between two classes of $(\alpha, \beta) - metrics$* , Diff. Geom. And its Applications, 27 (2009), 566-573.
- [13] H. S. Park and II-Yong Lee, *On Projectively flat Finsler spaces with $(\alpha, \beta) - metric$* , Comm. Korean Math. Soc., 14(2)(1999), 373-383.
- [14] H. S. Park and II-Yong Lee, *The Randers changes of Finsler spaces with $(\alpha, \beta) - metrics$ of Douglas type*, Comm. Korean Math. Soc., 38(3)(2001), 503-521.
- [15] H. S. Park and II-Yong Lee, *Projective changes between a Finsler space with $(\alpha, \beta) - metric$ and associated Riemannian metric*, Canad. J. Math., 60(2008), 443-456.
- [16] Pradeep Kumar, S.K. Narasimhamurthy, H.G. Nagaja and S.T. Aveesh, *On a special hypersurface of a Finsler space with $(\alpha, \beta) - metric$* , Tbilisi Mathematical Journal, 2(2009), 51-60.
- [17] Prasad C.M., and Dube O.P., *On T-Tensor and γ -curvature tensor of C3-like Finsler space*, Indian J. Pure Appl. Math., **23** (1992), 791-795.
- [18] B.N. Prasad, B.N Gupta and D.D. Singh, *Conformal transformation in Finsler spaces with $(\alpha, \beta) - metric$* , Indian J. Pure and Appl. Math., 18(4)(1961), 290-301.
- [19] M. Matsumoto and X. Wei, *Projective changes of Finsler spaces of constant curvature*, Publ. Math. Debreen, 44(1994), 175-181.
- [20] Rund H., *The differential Geometry of Finsler spaces*, Springer-Verlag, Berlin, 1959.